
Virginia COVID-19 Surveillance Data Update

April 28, 2022



US COVID-19: 7-Day Case Rate per 100,000, by State/Territory



	Cases in the Last 7 Days Per 100k Population
Virginia	79.4 (-8.9%)
U.S.	93.4 (+18.2%)
Vermont	284.9 (+2.1%)
New York New York City	249.2 (+8.3%) 209.7 (+7.0%)
District of Columbia	224.3 (+90.1%)

Our Neighbors

Rates Higher than Virginia

District of Columbia, **223.3** (+90.1%)

Maryland, **92.2** (+17.2%)


Rates Lower than Virginia:

Kentucky, **65.8** (-4.5%)

West Virginia, **50.6** (+39.4%)


Tennessee, **37.2** (+13.1%)


North Carolina, **28.5** (-60.3%)



- Omicron BA.2 sublineages account for 93.4% of US cases, as of 4/16/22
 - BA.2.12.1, an offshoot of BA.2, now accounts for 19% of US cases
 - BA.2.12.1 has been associated with rising cases and hospitalizations in central NY state

Central New York State Hospitalization Rates Rising







Compared to last week, **cases** increased to **44,308** (7-day MA) per day (+24.1%)

Hospitalizations decreased to **1,642** (7-day MA) per day (-6.6%)

Deaths decreased to **311** (7-day MA) per day (-18.2%)



Cases by Date of Illness for Virginia, Past 13 weeks



Gray shaded area illness may not have been reported yet

Confirmed Probable


Compared to last week, **cases increased** to 1,143 (7-day MA) from 1,096 per day (+4.3%)

Hospitalizations increased to **162** per day (7-day MA) (+22.7%)


Deaths decreased to 3 (-33.3%) (Date of Death)

* Death Data is usually Delayed in Reporting

Cases by Date Reported, All Reporting Timeline



COVID-19 in Virginia Hospitals




- Compared to last week hospitalizations increased to 162 (7-day MA) from 152 (+6.6%)
- Compared to last week ICU hospitalizations **decreased to 31** from 26 (-19.2%)

- Confirmed COVID-19 Patients Currently on Ventilator Support*^ ICU Hospitalizations (Confirmed)*
- CONFIRMED Hospitalizations*
- 7-Day Moving Average of Confirmed COVID-19 Hospitalizations

Medium

- If you are immunocompromised or <u>high</u> <u>risk</u> for severe disease
 - Talk to your healthcare provider about whether you need to wear a mask and take other precautions (e.g., testing)
 - Have a plan for rapid testing if needed (e.g., having home tests or access to testing)
 - Talk to your healthcare provider about whether you are a candidate for treatments like oral antivirals, PrEP, and monoclonal antibodies
- If you have household or social contact with someone at <u>high risk</u> for severe disease
 - consider self-testing to detect infection before contact
 - consider wearing a mask when indoors with them
- Stay up to date with COVID-19 vaccines and boosters
- Maintain improved ventilation throughout indoor spaces when possible
- Follow CDC recommendations for isolation and quarantine, including getting tested if you are exposed to COVID-19 or have symptoms of COVID-19




Time Period: COVID-19 Community Levels were calculated on Thu Apr 21 2022. New COVID-19 cases per 100,000 population (7-day total) are calculated using data from Thu Apr 14 2022 - Wed Apr 20 2022. New COVID-19 admissions per 100,000 population (7-day total) and Percent of inpatient beds occupied by COVID-19 patients (7-day average) are calculated using data from Wed Apr 13 2022 - Tue Apr 19 2022.

Key Trends

- There were 20 LTCF COVID-19 outbreaks reported in the past 30 days: 2 in Eastern, 6 in Central, 2 in Northwest, 4 in Northern, and 6 in Southwest (see figure top right).
- The number of reported staff cases slightly increased and resident cases in nursing homes slightly increased since the last reporting week (see figure bottom right).
 - For the reporting week ending April 24, 2022, 41 resident and 31 staff cases were reported to NHSN. Data for this reporting week are preliminary.
- For reporting week ending April 17, 2022, data reported by 281 nursing homes showed 89% of residents were fully vaccinated; data reported by 281 nursing homes showed 95% of staff were fully vaccinated (see figures bottom left). Of the nursing home residents eligible to receive an additional dose or booster, 81% of residents have received an additional dose or booster of COVID-19 vaccine.
 - Of the nursing home healthcare personnel eligible to receive an additional dose or booster, <u>46% of staff have received an additional dose or booster</u> of COVID-19 vaccine.


COVID-19 Booster Vaccination in Virginia Nursing Homes



Data were reported by 286 Virginia nursing homes into the National Healthcare Safety Network (NHSN) as of 4/25/2022 and are subject to change, including booster eligibility per <u>updated vaccine guidance</u>. In Virginia, 281 nursing homes reported resident vaccination data for reporting week ending 04/17/2022; 281 nursing homes reported staff vaccination data for reporting week ending 04/17/2022. For staff type definitions, refer to NHSN Table of Instructions.




COVID-19 Outbreaks in a Long-term Care Setting



Outbreaks reported from nursing homes, assisted living facilities, and multicare facilities to VDH with a confirmed or suspected etiologic agent of SARS-CoV-2. Data are from the Virginia Outbreak Surveillance System as of 04/25/2022; data are retrospectively updated and subject to change.

Nursing Home Resident and Staff COVID-19 Cases



Data are from NHSN as of 4/25/2022 and are subject to change. For reporting information, please refer to the NHSN data collection forms: residents, staff.

Metrics date: 4/25/2022


Central Far Southwest **Near Southwest** Northern Northwest Eastern New cases per 56.5 107.7 67.3 77.6 **157.3** 69.8 100k within the last 7 days % Positivity 7-day 5.6% 5.9% 8.5% 7.6% 9.7% 6.5% moving average **COVID-like ED visits 7.0** 4.7 10.3 9.6 9.6 6.0 rate per 100k

Burden	Level 0	Level 1	Level 2	Level 3	Level 4
New Cases	<10	10-49		50-100	>100
% Positivity	<3	3-5	5-8	8-10	>10
CLI ED Visits	<4		4-5.9		<u>></u> 6

Symbol	Trend
↑	Increasing
\	Decreasing
0	Fluctuating

Background

- Researchers observed increases in substance use and drug overdoses since the COVID-19 pandemic was declared a national emergency.
- Drug overdoses deaths accelerated during the COVID-19 pandemic. More than 93,000 drug overdose deaths were estimated in the United States in 2020, the highest number of overdose deaths ever recorded.
- People with current or previous SUD were 1.5 times more likely to have COVID-19 than those who did not.
- People with current or previous SUD were more likely to experience severe COVID-19 outcomes: hospitalization (41% versus 30%) and death (9.6% versus 6.6%).



Compounding COVID-19 Challenges for those with SUD and in Recovery

- Reduced social support, increased social isolation and increased stress
- Disruption of regular SUD treatment
- Increased rates of drug use as potential coping mechanism and use in isolation
- Decreased access to substance use treatment, harm reduction services, and emergency services

Opioid Users (heroin, fentanyl, morphine)

- Mechanism of Action: Causes slow breathing and decreased oxygen in blood and brain; increases risk for life-threatening overdose and damages the brain, heart, and lungs over time.
- Users are more susceptible to COVID-19 and at higher risk for more severe infections.
- COVID-19 hospitalization is 10.2 times more likely for those with opioid use disorder.


Stimulant Users (methamphetamine, cocaine, amphetamine)

- Mechanism of Action: Constricts the blood vessels and increases the risk for stroke, heart attacks, abnormal heart rhythm, seizures, and other conditions.
- Users are 6.5 times more likely to have COVID-19 and at higher risk for more severe infections
- COVID-19 hospitalization is 6.2 times more likely for those with stimulant use disorder

Homeless Population

- A high percentage of individuals with SUD experience homelessness
- People who experience homelessness and housing instability are at increased risk for COVID-19 due to higher rates of underlying health conditions and community spread in homeless shelters

People with a Lifetime or Recent SUD Diagnosis Are More Likely to Contract COVID-19 Than Those Without SUDs



Hospitalizations of Children Aged 5–11 Years with Laboratory-Confirmed COVID-19 | April 19, 2022, CDC MMWR **Summary:** COVID-NET data were analyzed to describe characteristics of COVID-19–associated hospitalizations among 1,475 U.S. children aged 5–11 from March 1, 2020–February 28, 2022, but mainly focusing on the period of early Omicron predominance.

Key Findings:


During the Omicron-predominant period:

- Cumulative hospitalization rates among unvaccinated children aged 5–11 years were 2.1 times as high as those among vaccinated children.
- **87% of hospitalized children** aged 5–11 years **were unvaccinated** (301 unvaccinated; 48 vaccinated).
- Among unvaccinated children, the largest proportion were Black (34%), followed by White (31%), and Hispanic (19%).
- There were no significant differences for severe outcomes by vaccination status (however the number of vaccinated children was very small; n = 48).

Across all variants:

- **32%** of hospitalized children aged 5–11 years had **severe COVID-19**
- The risk for severe COVID-19 among hospitalized children was significantly higher among those with diabetes (aRR = 2.5) and obesity (aRR = 1.2). Risk for severe disease was lower among children with asthma (aRR = 0.8), and those with immunocompromising conditions (aRR = 0.7)

Weekly Hospitalization Rates Among Children Aged 5-11 by Vaccination Status




Association of Short-term Air Pollution Exposure With SARS-CoV-2 Infection Among Young Adults in Sweden | April 20, 2022

Summary: A case-crossover study linking 425 participants who are part of the prospective BAMSE (Children, Allergy Milieu, Stockholm, Epidemiology [in Swedish]) birth cohort to the Swedish national infectious disease registry to identify cases with positive COVID-19 results from May 5, 2020, to March 31, 2021. Daily ambient air pollution levels at the individual residential address was calculated to estimate the association between short-term ambient air pollution and a positive PCR test.

Key Findings: Exposure to the following pollutants increased the risk of a positive SARS-CoV-2 PCR test:

- Exposure to **fine particulate matter** (PM_{2.5}) **two days before** a test was associated with a relative **increase of 6.8%** (95% CI, 2.1%-11.8%)
- Exposure to particulate matter (PM₁₀) two days before a test was associated with a relative increase of 6.9% (95% CI, 2.0%-12.1%)
- Exposure to **black carbon one day before** a test was associated with a relative **increase of 5.8%** (95% CI, 0.3%-11.6%)
- There was no observed association for exposure to nitrogen oxides
- The authors of the study speculate that increased levels of short-term air pollution plays a role in manifesting the disease (symptoms) for those who have been infected with the virus rather than contributing to the transmission of the virus (since median incubation period is 5 days).





Virginia Vaccination by Age

- ✓ **73.2%** (+0.4%) of the Total Population is Fully Vaccinated
- ✓ **38.9% (-0.8%)** of the Total Population is "Up-to-Date" with their Vaccinations
- ✓ **56.3%** (+0.2%) of the Eligible Population and **34.9%** (+1.3%) of Total Population Vaccinated with 3rd Dose/Booster
- ✓ 92.6% (+0.3%) of the Adult (18+) Population Vaccinated with at Least one Dose
- ✓ **58.4%** (+0.5%) of 5 to 17 year olds Vaccinated with at Least One Dose
- Green percent represents percent increase from two weeks prior


Second Booster Administrations Have Started


- Statewide, over 188,000 individuals have received their Second Booster
 - This accounts for about 6.3% of individuals with a First Booster
 - Average daily administrations of Second Boosters exceeded 6,000 last week






- The Black Population's Vaccination Rates are currently at 0.9x relative to the White Population's Rates
- The Parity between Latino and White Populations is much lower for Latinos when it comes to Third Dose/Booster Rates relative to First Dose (0.7x vs 1.2x)





Source: COVID-19 Vaccine Summary – Coronavirus (virginia.gov)

Percent of the Total Population Fully Vaccinated by Locality




2013 SRHP Isserman Classification	5 to 11	12 to 17	16 to 17	18 to 30	31 to 50	51 to 64	65+	Grand Total
Mixed Urban	41%	69%	73%	68%	69%	81%	88%	72%
Urban	38%	69%	75%	60%	72%	82%	86%	70%
Mixed Rural	25%	49%	56%	54%	60%	73%	82%	62%
Rural	17%	41%	47%	48%	54%	69%	78%	58%
Grand Total	34%	62%	67%	59%	67%	78%	84%	67%

Vaccination Rates by Region

Region Name	Region Name Fully Vaccinated	
Central	62.9%	38.3%
Eastern	58.9%	33.6%
Northern	73.7%	47.6%
Northwest	60.5%	36.4%
Southwest	53.5%	30.2%

- 21 out of 133 Localities have a fully vaccinated rate below 50%
- 15 out of 133 Localities have a fully vaccinated rate above 70%
- There is a disparity across Urban and Rural areas by Age Groups, with Rural Adolescents the Lowest Vaccinated group

Virginia and Neighbors: Vaccination Rates



	At Least One Dose*	Fully Vaccinated*	Booster Dose**
Nationwide	77.5% (+0.4%)	66.1% (+0.5%)	45.6% (+1.3%)
D.C.	95.0% (+0.0%)	73.9% (+0.8%)	37.9% (+3.8%)
Kentucky	66.1% (+0.3%)	57.3% (+0.4%)	44.1% (+0.9%)
Maryland	86.3% (+-0.2%)	75.5% (+-0.4%)	50.6% (+1.2%)
North Carolina	84.0% (+0.6%)	60.9% (+1%)	26.4% (+1.1%)
Tennessee	62.1% (+0.3%)	54.4% (+0.4%)	43.8% (+1.4%)
Virginia**	85.5% (+0.4%)	73.1% (+0.4%)	47.1% (+0.9%)
West Virginia	64.9% (+0.3%)	57.5% (+0.3%)	45.6% (+1.3%)

*Total population, includes out-of-state vaccinations

**Percent of fully vaccinated people with a booster dose

***Differs from previous slide because all vaccination
sources (e.g., federal) are included

**** Green percent represents percent increase from
three weeks prior